

ICPSR 33444

**Center for Education and Drug
Abuse Research (CEDAR):
Etiological and Prospective Family
Study in Southwestern
Pennsylvania, Baseline Data,
1990-2011**

Ralph E. Tarter
University of Pittsburgh

Nursing Assessment, Variable Description and
Frequencies

Inter-university Consortium for
Political and Social Research
P.O. Box 1248
Ann Arbor, Michigan 48106
www.icpsr.umich.edu

Terms of Use

The terms of use for this study can be found at:
<http://www.icpsr.umich.edu/icpsrweb/ICPSR/studies/33444/terms>

Information about Copyrighted Content

Some instruments administered as part of this study may contain in whole or substantially in part contents from copyrighted instruments. Reproductions of the instruments are provided as documentation for the analysis of the data associated with this collection. Restrictions on "fair use" apply to all copyrighted content. More information about the reproduction of copyrighted works by educators and librarians is available from the United States Copyright Office.

NOTICE

WARNING CONCERNING COPYRIGHT RESTRICTIONS

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.

Data Completeness Report

Notes: (1) Variables are individually listed only if they have greater than 5% missing data. These variables are listed under the appropriate percentage category in the order in which they appear in the data file. (2) The Data Completeness Report only captures information about system missing or other values that are declared missing. Codes that have a label implying that they are missing but that are not declared missing values are not reflected in this report. Data users should consult the codebook for more specific information about missing values. (3) Some variables that have 100% missing data may have been blanked by ICPSR to protect respondent confidentiality. Data users should consult the codebook for more specific information about blanked variables. (4) Data do not contain skip patterns or skip patterns are not reflected in the data as coded.

Table 1: Distribution of Variables by Percentage of Missing Values--Nursing Assessment

Variable Name and Label (Total Cases = 773)	Percent of Cases with Missing Values
33.3% (4 of 12 variables)	have 0% Missing Values
41.7% (5 of 12 variables)	have 0% - 1% Missing Values
25.0% (3 of 12 variables)	have 1% - 3% Missing Values
0.0% (0 of 12 variables)	have 3% - 5% Missing Values
0.0% (0 of 12 variables)	have 5% - 10% Missing Values
0.0% (0 of 12 variables)	have 10% - 20% Missing Values
0.0% (0 of 12 variables)	have 20% - 40% Missing Values
0.0% (0 of 12 variables)	have 40% - 99% Missing Values
0.0% (0 of 12 variables)	have 100% missing values

ICPSR 33444

Center for Education and Drug Abuse Research (CEDAR): Etiological and Prospective Family Study in Southwestern Pennsylvania, Baseline Data, 1990-2011 [United States]

Variable Description and Frequencies

Note: Frequencies displayed for the variables are not weighted. They are purely descriptive and may not be representative of the study population. Please review any sampling or weighting information available with the study.

Summary statistics (minimum, maximum, mean, median, and standard deviation) may not be available for every variable in the codebook. Conversely, a listing of frequencies in table format may not be present for every variable in the codebook either. However, all variables in the dataset are present and display sufficient information about each variable. These decisions are made intentionally and are at the discretion of the archive producing this codebook.

Center for Education and Drug Abuse Research (CEDAR): Etiological and Prospective Family Study in Southwestern Pennsylvania, Baseline Data, 1990-2011 [United States]

Nursing Assessment

ID Subject ID

Location: 1-11 (width: 11; decimal: 0)
Variable Type: character
Based upon 773 valid cases out of 773 total cases.

ASSID Assessment ID

Location: 12-22 (width: 11; decimal: 0)
Variable Type: character
Based upon 773 valid cases out of 773 total cases.

VISIT Visit

Location: 23-24 (width: 2; decimal: 0)
Variable Type: numeric

<i>Value</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	773	100.0 %

Based upon 773 valid cases out of 773 total cases.

- Mean: 1.00
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 1.00
- Standard Deviation: 0.00

NA1 1. Height (in.)

Location: 25-33 (width: 9; decimal: 1)
Variable Type: numeric
Range of Missing Values (M): -2.0

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
48.5	-	1	0.1 %
50.0	-	4	0.5 %
50.5	-	3	0.4 %
51.0	-	5	0.6 %
51.3	-	3	0.4 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
51.5	-	1	0.1 %
51.8	-	1	0.1 %
52.0	-	4	0.5 %
52.3	-	6	0.8 %
52.5	-	7	0.9 %
52.8	-	3	0.4 %
53.0	-	14	1.8 %
53.3	-	5	0.6 %
53.5	-	11	1.4 %
53.8	-	5	0.6 %
54.0	-	21	2.7 %
54.3	-	10	1.3 %
54.5	-	20	2.6 %
54.8	-	10	1.3 %
55.0	-	15	1.9 %
55.2	-	1	0.1 %
55.3	-	14	1.8 %
55.5	-	24	3.1 %
55.8	-	13	1.7 %
56.0	-	27	3.5 %
56.2	-	1	0.1 %
56.3	-	10	1.3 %
56.5	-	19	2.5 %
56.8	-	14	1.8 %
57.0	-	26	3.4 %
57.3	-	18	2.3 %
57.5	-	21	2.7 %
57.8	-	14	1.8 %
58.0	-	26	3.4 %
58.3	-	15	1.9 %
58.5	-	23	3.0 %
58.8	-	18	2.3 %
59.0	-	30	3.9 %
59.3	-	19	2.5 %
59.5	-	21	2.7 %
59.8	-	8	1.0 %
60.0	-	24	3.1 %
60.1	-	2	0.3 %
60.3	-	8	1.0 %
60.5	-	22	2.8 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
60.6	-	1	0.1 %
60.8	-	11	1.4 %
61.0	-	18	2.3 %
61.3	-	8	1.0 %
61.5	-	13	1.7 %
61.8	-	14	1.8 %
62.0	-	12	1.6 %
62.3	-	7	0.9 %
62.5	-	13	1.7 %
62.8	-	7	0.9 %
63.0	-	19	2.5 %
63.3	-	8	1.0 %
63.5	-	7	0.9 %
63.8	-	4	0.5 %
64.0	-	10	1.3 %
64.3	-	4	0.5 %
64.5	-	8	1.0 %
64.8	-	4	0.5 %
65.0	-	5	0.6 %
65.3	-	1	0.1 %
65.5	-	4	0.5 %
65.8	-	1	0.1 %
66.0	-	4	0.5 %
66.3	-	2	0.3 %
66.5	-	3	0.4 %
66.8	-	2	0.3 %
67.0	-	2	0.3 %
67.3	-	2	0.3 %
67.7	-	1	0.1 %
68.0	-	1	0.1 %
68.3	-	1	0.1 %
69.0	-	1	0.1 %
69.3	-	1	0.1 %
69.5	-	3	0.4 %
-2.0 (M)	Missing, N/A	4	0.5 %

Based upon 769 valid cases out of 773 total cases.

- Mean: 58.42
- Median: 58.30
- Mode: 59.00

- Study 33444 -

- Minimum: 48.50
- Maximum: 69.50
- Standard Deviation: 3.67

NA2 2. Weight (lbs.)

Location: 34-42 (width: 9; decimal: 1)
 Variable Type: numeric
 Range of Missing Values (M): -2

<i>Value</i>	<i>Label</i>
-2 (M)	Missing, N/A

Based upon 772 valid cases out of 773 total cases.

- Mean: 98.42
- Minimum: 50.50
- Maximum: 236.00
- Standard Deviation: 28.67

NA3A 3a. Blood Pressure: Systolic

Location: 43-45 (width: 3; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
71	-	1	0.1 %
72	-	4	0.5 %
77	-	1	0.1 %
78	-	9	1.2 %
80	-	35	4.5 %
82	-	17	2.2 %
84	-	13	1.7 %
86	-	12	1.6 %
87	-	1	0.1 %
88	-	51	6.6 %
90	-	114	14.7 %
91	-	1	0.1 %
92	-	36	4.7 %
93	-	1	0.1 %
94	-	22	2.8 %
95	-	4	0.5 %
96	-	21	2.7 %
98	-	70	9.1 %
99	-	2	0.3 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
100	-	104	13.5 %
102	-	21	2.7 %
103	-	2	0.3 %
104	-	19	2.5 %
105	-	2	0.3 %
106	-	9	1.2 %
107	-	6	0.8 %
108	-	22	2.8 %
109	-	3	0.4 %
110	-	50	6.5 %
111	-	1	0.1 %
112	-	4	0.5 %
113	-	4	0.5 %
114	-	4	0.5 %
115	-	8	1.0 %
116	-	5	0.6 %
117	-	3	0.4 %
118	-	13	1.7 %
119	-	4	0.5 %
120	-	32	4.1 %
121	-	1	0.1 %
122	-	3	0.4 %
123	-	3	0.4 %
124	-	2	0.3 %
125	-	1	0.1 %
126	-	1	0.1 %
127	-	1	0.1 %
128	-	6	0.8 %
129	-	3	0.4 %
130	-	6	0.8 %
133	-	3	0.4 %
135	-	1	0.1 %
138	-	2	0.3 %
139	-	1	0.1 %
140	-	1	0.1 %
144	-	1	0.1 %
150	-	1	0.1 %
159	-	1	0.1 %
161	-	1	0.1 %
-2 (M)	Missing, N/A	3	0.4 %

- Study 33444 -

Based upon 770 valid cases out of 773 total cases.

- Mean: 99.13
- Median: 98.00
- Mode: 90.00
- Minimum: 71.00
- Maximum: 161.00
- Standard Deviation: 13.00

NA3B

3b. Blood Pressure: Diastolic

Location: 46-48 (width: 3; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
40	-	2	0.3 %
42	-	1	0.1 %
44	-	1	0.1 %
45	-	1	0.1 %
48	-	8	1.0 %
50	-	34	4.4 %
51	-	1	0.1 %
52	-	31	4.0 %
53	-	1	0.1 %
54	-	11	1.4 %
55	-	2	0.3 %
56	-	12	1.6 %
57	-	2	0.3 %
58	-	87	11.3 %
59	-	4	0.5 %
60	-	202	26.1 %
61	-	3	0.4 %
62	-	50	6.5 %
63	-	4	0.5 %
64	-	26	3.4 %
65	-	5	0.6 %
66	-	13	1.7 %
67	-	4	0.5 %
68	-	61	7.9 %
69	-	4	0.5 %
70	-	77	10.0 %
71	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
72	-	20	2.6 %
73	-	4	0.5 %
74	-	8	1.0 %
75	-	2	0.3 %
76	-	2	0.3 %
77	-	2	0.3 %
78	-	19	2.5 %
79	-	1	0.1 %
80	-	37	4.8 %
81	-	1	0.1 %
82	-	8	1.0 %
84	-	5	0.6 %
86	-	2	0.3 %
88	-	3	0.4 %
90	-	3	0.4 %
91	-	1	0.1 %
93	-	1	0.1 %
98	-	1	0.1 %
105	-	1	0.1 %
-2 (M)	Missing, N/A	4	0.5 %

Based upon 769 valid cases out of 773 total cases.

- Mean: 63.70
- Median: 60.00
- Mode: 60.00
- Minimum: 40.00
- Maximum: 105.00
- Standard Deviation: 8.79

NA4

4. Pulse

Location: 49-51 (width: 3; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
54	-	1	0.1 %
56	-	3	0.4 %
58	-	1	0.1 %
60	-	25	3.2 %
62	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
63	-	3	0.4 %
64	-	33	4.3 %
65	-	3	0.4 %
66	-	1	0.1 %
68	-	65	8.4 %
69	-	2	0.3 %
70	-	3	0.4 %
71	-	7	0.9 %
72	-	97	12.5 %
73	-	4	0.5 %
74	-	6	0.8 %
75	-	3	0.4 %
76	-	74	9.6 %
77	-	3	0.4 %
78	-	14	1.8 %
79	-	3	0.4 %
80	-	111	14.4 %
81	-	4	0.5 %
82	-	10	1.3 %
84	-	66	8.5 %
85	-	2	0.3 %
86	-	5	0.6 %
87	-	2	0.3 %
88	-	93	12.0 %
89	-	4	0.5 %
90	-	5	0.6 %
91	-	1	0.1 %
92	-	36	4.7 %
93	-	1	0.1 %
94	-	2	0.3 %
96	-	32	4.1 %
97	-	1	0.1 %
98	-	1	0.1 %
99	-	1	0.1 %
100	-	22	2.8 %
102	-	2	0.3 %
104	-	4	0.5 %
106	-	1	0.1 %
108	-	4	0.5 %
110	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
112	-	2	0.3 %
116	-	2	0.3 %
121	-	1	0.1 %
128	-	1	0.1 %
-2 (M)	Missing, N/A	4	0.5 %

Based upon 769 valid cases out of 773 total cases.

- Mean: 79.78
- Median: 80.00
- Mode: 80.00
- Minimum: 54.00
- Maximum: 128.00
- Standard Deviation: 10.85

NA5 5. Respirations

Location: 52-54 (width: 3; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
12	-	21	2.7 %
14	-	281	36.4 %
16	-	135	17.5 %
17	-	3	0.4 %
18	-	115	14.9 %
20	-	189	24.5 %
22	-	5	0.6 %
24	-	17	2.2 %
25	-	7	0.9 %

Based upon 773 valid cases out of 773 total cases.

- Mean: 16.74
- Median: 16.00
- Mode: 14.00
- Minimum: 12.00
- Maximum: 25.00
- Standard Deviation: 2.87

NA6 6. Temperature

Location: 55-63 (width: 9; decimal: 1)

- Study 33444 -

Variable Type: numeric

Range of Missing Values (M): -2.0

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
90.0	-	1	0.1 %
95.2	-	1	0.1 %
95.7	-	1	0.1 %
95.9	-	1	0.1 %
96.0	-	1	0.1 %
96.1	-	2	0.3 %
96.2	-	4	0.5 %
96.4	-	5	0.6 %
96.5	-	1	0.1 %
96.6	-	4	0.5 %
96.7	-	4	0.5 %
96.8	-	5	0.6 %
96.9	-	7	0.9 %
97.0	-	4	0.5 %
97.1	-	11	1.4 %
97.2	-	17	2.2 %
97.3	-	8	1.0 %
97.4	-	14	1.8 %
97.5	-	18	2.3 %
97.6	-	26	3.4 %
97.7	-	21	2.7 %
97.8	-	55	7.1 %
97.9	-	32	4.1 %
98.0	-	43	5.6 %
98.1	-	40	5.2 %
98.2	-	41	5.3 %
98.3	-	27	3.5 %
98.4	-	36	4.7 %
98.5	-	35	4.5 %
98.6	-	46	6.0 %
98.7	-	33	4.3 %
98.8	-	46	6.0 %
98.9	-	48	6.2 %
99.0	-	29	3.8 %
99.1	-	18	2.3 %
99.2	-	22	2.8 %
99.3	-	15	1.9 %
99.4	-	17	2.2 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
99.5	-	9	1.2 %
99.6	-	5	0.6 %
99.7	-	1	0.1 %
99.8	-	4	0.5 %
99.9	-	3	0.4 %
100.0	-	2	0.3 %
100.1	-	1	0.1 %
100.6	-	1	0.1 %
-2.0 (M)	Missing, N/A	8	1.0 %

Based upon 765 valid cases out of 773 total cases.

- Mean: 98.26
- Median: 98.30
- Mode: 97.80
- Minimum: 90.00
- Maximum: 100.60
- Standard Deviation: 0.80

NA7

7. Weber

Location: 64-66 (width: 3; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	DNL	670	86.7 %
2	R	60	7.8 %
3	L	25	3.2 %
-2 (M)	Missing, N/A	18	2.3 %

Based upon 755 valid cases out of 773 total cases.

- Mean: 1.15
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 3.00
- Standard Deviation: 0.44

NA8

8. Rinne

Location: 67-69 (width: 3; decimal: 0)
 Variable Type: numeric

- Study 33444 -

Range of Missing Values (M): -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	AC > BC	754	97.5 %
2	AC < BC	5	0.6 %
-2 (M)	Missing, N/A	14	1.8 %

Based upon 759 valid cases out of 773 total cases.

- Mean: 1.01
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 2.00
- Standard Deviation: 0.08