

ICPSR 33444

**Center for Education and Drug  
Abuse Research (CEDAR):  
Etiological and Prospective Family  
Study in Southwestern  
Pennsylvania, Baseline Data,  
1990-2011**

Ralph E. Tarter  
*University of Pittsburgh*

Psychiatric History Marker for DSM-IV - Index  
Case, Data Collection Instrument

Inter-university Consortium for  
Political and Social Research  
P.O. Box 1248  
Ann Arbor, Michigan 48106  
[www.icpsr.umich.edu](http://www.icpsr.umich.edu)

# Terms of Use

The terms of use for this study can be found at:  
<http://www.icpsr.umich.edu/icpsrweb/ICPSR/studies/33444/terms>

## Information about Copyrighted Content

Some instruments administered as part of this study may contain in whole or substantially in part contents from copyrighted instruments. Reproductions of the instruments are provided as documentation for the analysis of the data associated with this collection. Restrictions on "fair use" apply to all copyrighted content. More information about the reproduction of copyrighted works by educators and librarians is available from the United States Copyright Office.

### NOTICE

#### WARNING CONCERNING COPYRIGHT RESTRICTIONS

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.

## PSYCHIATRIC HISTORY MARKER DSM-IV

File name: CEDAR\_psychiatric\_history\_DSMIV\_dad.sav  
CEDAR\_psychiatric\_history\_DSMIV\_ic.sav  
CEDAR\_psychiatric\_history\_DSMIV\_mom.sav

Done by: Father, Mother, Index Case

Source: Center for Education and Drug Abuse Research (CEDAR) (unpublished). University of Pittsburgh.

---

<u>COLUMN</u>	<u>QUESTION TEXT</u>
ID	SUBJECT ID:
ASSID	ASSESSMENT ID:
VISIT	VISIT NUMBER:
	<b>PSYCHIATRIC DISORDER</b>
D401	DIAGNOSIS #
D402	1 = DEFINITE    2 = PROBABLE
	DSM-IV
D403A	TEXT
D407	DSM-IV CODE:
	ICD-10
D408	TEXT
D409	ICD-10 CODE:
	<b>II. CHRONOLOGY (LIFETIME)</b>
D410	1) Age of onset ____ (Years)
D411	2) Age of offset ____ (Years)
	<b>III. COURSE SINCE THE ONSET OF THE SYNDROME</b>
D412	1) Continuous or single episodes                      DK = -1    Yes = 1    No = 0
D413	2) Fluctuating without clear-cut episodes              DK = -1    Yes = 1    No = 0
D414	3 a) Clear-cut episodes                                      DK = -1    Yes = 1    No = 0
D415	b) Number of episodes                                      _____

<u>COLUMN</u>	<u>COLUMN</u>	<b>IV. CONTRIBUTING FACTORS</b>
<b>CURRENT</b>	<b>PAST</b>	
D416	D417	1) Age of onset
D418	D419	2) Age of offset
D420D	D421D	3) Duration in days

**CURRENT PAST**

D422	D423	B. DRUG OR ALCOHOL	<b>Responses for questions B, C, D</b> 1 = None 2 = Antecedent or Consequent 3 = Exacerbation 4 = Maintenance 5 = Alleviate
D424	D425	C. MEDICATION	
D426	D427	D. PHYSICAL ILLNESS	
D428	D429	E. ENVIRONMENTAL STRESSORS	
		<b>V. TREATMENT</b>	<b>-1 = DK    0 = NO    1 = YES</b>
D430	D431	A. 1) Inpatient	
D432	D433	2) Outpatient	
D434	D435	3) Medication	
		<b>B. MODALITY</b>	
D436	D437	1) Individual therapy	
D438	D439	2) Group Therapy	
D440	D441	3) Family Therapy	
D442	D443	4) Specific Behavioral Technique	
D444	D445	5) Vocational training	
D446	D447	6) Neuroleptics	
D448	D449	7) Lithium	
D450	D451	8) Antidepressants	
D452	D453	9) Minor Tranquilizers	
D454	D455	10) Stimulants	
D456	D457	11) Anticonvulsants	
D458	D459	12) Anti-Parkinson's medication	
D460	D461	13) Other/Unspecified medication	
D462	D463	14) ECT	
D464	D465	15) Other/Unspecified modality	
		<b>VI. SOCIAL FUNCTIONING AFFECTED</b>	
D466	D467	1) Family	
D468	D469	2) Peers	
D470	D471	3) Work/school	
D472	D473	4) Community	
D474		VII. Global Assessment Scale (when applicable) _____ (0 -100) <b>See below</b>	

---

**GLOBAL ASSESSMENT OF FUNCTIONING (GAF) SCALE  
DSM-IV**

Consider psychological, social and occupational functioning on a hypothetical continuum of mental health-illness. Do not include impairment in functioning due to physical (or environmental) limitations.

**CODE FOR D474** (Note: Use intermediate codes when appropriate, e.g., 45, 68.72.)

100 Superior functioning in a wide range of activities, life's problems never seem to get out of hand, is sought out by others because of his or her many positive qualities. No symptoms  
|  
91

90 Absent or minimal symptoms (e.g., mild anxiety before an exam), good functioning in all areas,  
| interested and involved in a wide range of activities, socially effective, generally satisfied with life,  
81 no more than everyday problems or concerns (e.g., an occasional argument with family members).

- 80 | If symptoms are present, they are transient and expected reactions to psychosocial  
71 | stressors (e.g., difficulty concentration after family argument); no more than slight impairment  
in social, occupational, or school functioning (e.g., temporarily falling behind in schoolwork)
- 70 | Some mild symptoms (e.g., depressed mood and mild insomnia) OR some difficulty in social,  
61 | occupational, or school functioning (e.g., occasional truancy, or theft within the household).  
but generally functioning pretty well, has some meaningful interpersonal relationships.
- 60 | Moderate symptoms (e.g., flat affect and circumstantial speech, occasional panic attacks) OR  
51 | moderate difficulty in social, occupational, or school functioning (e.g., few friends, conflicts  
with peers or co-workers)
- 50 | Serious symptoms (e.g., suicidal ideation, severe obsessive rituals, frequent shoplifting) OR any  
41 | serious impairment in social, occupational, or school functioning (e.g., no friends, unable to  
keep a job).
- 40 | Some impairment in reality testing or communication (e.g., speech is at times illogical, obscure  
| or irrelevant) OR major impairment in several areas, such as work or school, family  
31 | relations, judgment, thinking or mood (e.g., depressed man avoids friends, neglects family, and is  
unable to work, child frequently beats up younger children, is defiant at home, and is failing in school)
- 30 | Behavior is considerably influenced by delusion or hallucinations OR serious impairment in  
| communication or judgment (e.g., sometimes incoherent, acts grossly inappropriately, suicidal  
21 | preoccupation) OR inability to function in almost all areas (e.g., stays in bed all day; no job,  
home, or friends)
- 20 | Some danger of hurting self or others (e.g., suicide attempts without clear expectation of death;  
| frequently violent; manic excitement) OR occasionally fails to maintain minimal personal hygiene  
11 | (e.g., smears feces) OR gross impairment in communication (e.g., largely incoherent or mute)
- 10 | Persistent danger of severely hurting self or others (e.g., recurrent violence) OR persistent  
| inability to maintain minimal personal hygiene OR serious suicidal act with clear expectation of death  
01
- 0 | Inadequate information