

ICPSR 33444

**Center for Education and Drug
Abuse Research (CEDAR):
Etiological and Prospective Family
Study in Southwestern
Pennsylvania, Baseline Data,
1990-2011**

Ralph E. Tarter
University of Pittsburgh

Psychiatric Treatment History Questionnaire -
Father, Variable Description and Frequencies

Inter-university Consortium for
Political and Social Research
P.O. Box 1248
Ann Arbor, Michigan 48106
www.icpsr.umich.edu

Terms of Use

The terms of use for this study can be found at:
<http://www.icpsr.umich.edu/icpsrweb/ICPSR/studies/33444/terms>

Information about Copyrighted Content

Some instruments administered as part of this study may contain in whole or substantially in part contents from copyrighted instruments. Reproductions of the instruments are provided as documentation for the analysis of the data associated with this collection. Restrictions on "fair use" apply to all copyrighted content. More information about the reproduction of copyrighted works by educators and librarians is available from the United States Copyright Office.

NOTICE

WARNING CONCERNING COPYRIGHT RESTRICTIONS

The copyright law of the United States (Title 17, United States Code) governs the making of photocopies or other reproductions of copyrighted material. Under certain conditions specified in the law, libraries and archives are authorized to furnish a photocopy or other reproduction. One of these specified conditions is that the photocopy or reproduction is not to be "used for any purpose other than private study, scholarship, or research." If a user makes a request for, or later uses, a photocopy or reproduction for purposes in excess of "fair use," that user may be liable for copyright infringement.

ICPSR PROCESSING NOTES FOR #33444

*Center for Education and Drug Abuse Research (CEDAR): Etiological and Prospective Family Study in Southwestern Pennsylvania, Baseline Data, 1990-2011 [United States]
Part 38: Psychiatric Treatment History Questionnaire - Father*

- 1) For confidentiality reasons, ICPSR masked information in some character variables in the data. Changed text is denoted by square brackets, [].
- 2) Valid values in character variables may not be displayed in the frequency tables. Refer to the frequencies for missing data and/or the statement below the table for information on how many cases contain valid values.
- 3) Due to the means by which SAS sets the length of a variable based on the first occurrence of the variable, ICPSR removed value labels from the following long character variables for the SAS transport files and the SAS setup files.

PCU02

PCU07

PCU12

- 4) Some continuous variables have high or low values that may fall outside the valid range for that variable. These values were not labeled because they may be valid. However, users of the data should be aware of the possibility that these values may not be valid.

Data Completeness Report

Notes: (1) Variables are individually listed only if they have greater than 5% missing data. These variables are listed under the appropriate percentage category in the order in which they appear in the data file. (2) The Data Completeness Report only captures information about system missing or other values that are declared missing. Codes that have a label implying that they are missing but that are not declared missing values are not reflected in this report. Data users should consult the codebook for more specific information about missing values. (3) Some variables that have 100% missing data may have been blanked by ICPSR to protect respondent confidentiality. Data users should consult the codebook for more specific information about blanked variables. (4) Data do not contain skip patterns or skip patterns are not reflected in the data as coded.

Table 1: Distribution of Variables by Percentage of Missing Values--Psychiatric Treatment History Questionnaire - Father

Variable Name and Label (Total Cases = 715)	Percent of Cases with Missing Values
13.6% (12 of 88 variables)	have 0% Missing Values
3.4% (3 of 88 variables)	have 0% - 1% Missing Values
0.0% (0 of 88 variables)	have 1% - 3% Missing Values
0.0% (0 of 88 variables)	have 3% - 5% Missing Values
0.0% (0 of 88 variables)	have 5% - 10% Missing Values
0.0% (0 of 88 variables)	have 10% - 20% Missing Values
0.0% (0 of 88 variables)	have 20% - 40% Missing Values
83.0% (73 of 88 variables)	have 40% - 99% Missing Values
PCA21	How old were you the first time you went for help because of emotional problems? 73.6%
PCA31	How old were you the last time you went for help because of emotional problems? 73.6%
PCA41	How many times were you in treatment as an outpatient because of emotional problems? 75.2%
PCA51	How many times were you in treatment as an inpatient because of emotional problems? 92.2%
PCA22	How old were you the first time you went for help because of drinking problems? 85.0%
PCA32	How old were you the last time you went for help because of drinking problems? 85.0%
PCA42	How many times were you in treatment as an outpatient because of drinking problems? 88.4%
PCA52	How many times were you in treatment as an inpatient because of drinking problems? 88.3%
PCA23	How old were you the first time you went for help because of drug problems? 78.5%
PCA33	How old were you the last time you went for help because of drug problems? 78.5%
PCA43	How many times were you in treatment as an outpatient because of drug problems? 83.6%

Table 1: Distribution of Variables by Percentage of Missing Values--Psychiatric Treatment History Questionnaire - Father

Variable Name and Label (Total Cases = 715)		Percent of Cases with Missing Values
PCA53	How many times were you in treatment as an inpatient because of drug problems?	81.5%
PCA81	Whom did you see most frequently as an outpatient for Emotional Problems?	77.1%
PCA82	Whom did you see most frequently as an outpatient for Drinking Problems?	91.3%
PCA83	Whom did you see most frequently as an outpatient for Drug Problems?	87.4%
PCA84	Whom did you see most frequently as an Inpatient for Emotional Problems?	93.3%
PCA85	Whom did you see most frequently as an Inpatient for Drinking Problems?	88.7%
PCA86	Whom did you see most frequently as an Inpatient for Drug Problems?	82.7%
PCOE9	How long did you go to this person as an Outpatient for Emotional Problems?	77.1%
PCOG9	How long did you go to this person as an Outpatient for Drinking Problems?	91.3%
PCOR9	How long did you go to this person as an Outpatient for Drug Problems?	87.6%
PCIE9	How long did you go to this person as an Inpatient for Emotional Problems?	93.3%
PCIG9	How long did you go to this person as an Inpatient for Drinking Problems?	88.7%
PCIR9	How long did you go to this person as an Inpatient for Drug Problems?	82.7%
PC101	What type of treatment did you have because of Emotional Problems - Outpatient Psychotherapy?	76.9%
PC105	What type of treatment did you have because of Drinking Problems - Outpatient Psychotherapy?	91.3%
PC111	What type of treatment did you have because of Drug Problems - Outpatient Psychotherapy?	87.4%
PC102	What type of treatment did you have because of Emotional Problems - Outpatient ECT?	77.1%
PC106	What type of treatment did you have because of Drinking Problems - Outpatient ECT?	91.3%
PC112	What type of treatment did you have because of Drug Problems - Outpatient ECT?	87.4%
PC103	What type of treatment did you have because of Emotional Problems - Outpatient Medication?	77.1%
PC107	What type of treatment did you have because of Drinking Problems - Outpatient Medication?	91.3%
PC113	What type of treatment did you have because of Drug Problems - Outpatient Medication?	87.4%

Table 1: Distribution of Variables by Percentage of Missing Values--Psychiatric Treatment History Questionnaire - Father

Variable Name and Label (Total Cases = 715)	Percent of Cases with Missing Values
PC108 What type of treatment did you have because of Drinking Problems - Outpatient Detoxification?	91.3%
PC114 What type of treatment did you have because of Drug Problems - Outpatient Detoxification?	87.4%
PC109 What type of treatment did you have because of Drinking Problems - Outpatient Rehabilitation?	91.3%
PC115 What type of treatment did you have because of Drug Problems - Outpatient Rehabilitation?	87.4%
PC104 What type of treatment did you have because of Emotional Problems - Other Outpatient?	77.2%
PC110 What type of treatment did you have because of Drinking Problems - Other Outpatient	91.3%
PC116 What type of treatment did you have because of Drug Problems - Other Outpatient	87.4%
PC117 What type of treatment did you have because of Emotional Problems - Inpatient Psychotherapy?	92.9%
PC123 What type of treatment did you have because of Drinking Problems - Inpatient Psychotherapy?	88.7%
PC129 What type of treatment did you have because of Drug Problems - Inpatient Psychotherapy?	82.8%
PC118 What type of treatment did you have because of Emotional Problems - Inpatient ECT?	92.9%
PC124 What type of treatment did you have because of Drinking Problems - Inpatient ECT?	88.7%
PC130 What type of treatment did you have because of Drug Problems - Inpatient ECT?	82.8%
PC119 What type of treatment did you have because of Emotional Problems - Inpatient Medication?	92.9%
PC125 What type of treatment did you have because of Drinking Problems - Inpatient Medication?	88.7%
PC131 What type of treatment did you have because of Drug Problems - Inpatient Medication?	82.8%
PC120 What type of treatment did you have because of Emotional Problems - Inpatient Detoxification?	92.9%
PC126 What type of treatment did you have because of Drinking Problems - Inpatient Detoxification?	88.7%
PC132 What type of treatment did you have because of Drug Problems - Inpatient Detoxification?	82.8%
PC121 What type of treatment did you have because of Emotional Problems - Inpatient Rehabilitation?	92.9%
PC127 What type of treatment did you have because of Drinking Problems - Inpatient Rehabilitation?	88.7%

Table 1: Distribution of Variables by Percentage of Missing Values--Psychiatric Treatment History Questionnaire - Father

Variable Name and Label (Total Cases = 715)	Percent of Cases with Missing Values
PC133 What type of treatment did you have because of Drug Problems - Inpatient Rehabilitation?	82.7%
PC122 What type of treatment did you have because of Emotional Problems - Other Inpatient?	92.9%
PC128 What type of treatment did you have because of Drinking Problems - Other Inpatient?	88.7%
PC134 What type of treatment did you have because of Drug Problems - Other Inpatient?	82.9%
PCU02 Worst time you needed professional help because of emotional problems and you did not get it: 2) What was the reason?	90.5%
PCU03 Worst time you needed professional help because of emotional problems and you did not get it: 3) How old were you when you first needed help?	89.9%
PCU04 Worst time you needed professional help because of emotional problems and you did not get it: 4) How old were you the last time you needed help?	89.8%
PCU05 Worst time you needed professional help because of emotional problems and you did not get it: 5) How many times did you feel you needed help?	90.3%
PCU06 Worst time you needed professional help because of emotional problems and you did not get it: 6) During this period for how long (# days) did you need help but not get it	89.9%
PCU07 Worst time you needed professional help because of drinking problems and you did not get it: 2) What was the reason?	92.4%
PCU08 Worst time you needed professional help because of drinking problems and you did not get it: 3) How old were you when you first needed help?	91.7%
PCU09 Worst time you needed professional help because of drinking problems and you did not get it: 4) How old were you the last time you needed help?	91.7%
PCU10 Worst time you needed professional help because of drinking problems and you did not get it: 5) How many times did you feel you needed help?	92.2%
PCU11 Worst time you needed professional help because of drinking problems and you did not get it: 6) During this period for how long (# days) did you need help but not get it?	91.9%
PCU12 Worst time you needed professional help because of drug problems and you did not get it: 2) What was the reason?	90.5%
PCU14 Worst time you needed professional help because of drug problems and you did not get it: 4) How old were you when you last needed help?	89.7%
PCU13 Worst time you needed professional help because of drug problems and you did not get it: 3) How old were you the first time you needed help?	89.7%

Table 1: Distribution of Variables by Percentage of Missing Values--Psychiatric Treatment History Questionnaire - Father

Variable Name and Label (Total Cases = 715)	Percent of Cases with Missing Values
PCU15	Worst time you needed professional help because of drug problems and you did not get it: 5) How many times did you feel you needed help? 89.9%
PCU16	Worst time you needed professional help because of drug problems and you did not get it: 6) During this period for how long (# days) did you need help but not get it? 89.5%
0.0% (0 of 88 variables)	have 100% missing values

ICPSR 33444

Center for Education and Drug Abuse Research (CEDAR): Etiological and Prospective Family Study in Southwestern Pennsylvania, Baseline Data, 1990-2011 [United States]

Variable Description and Frequencies

Note: Frequencies displayed for the variables are not weighted. They are purely descriptive and may not be representative of the study population. Please review any sampling or weighting information available with the study.

Summary statistics (minimum, maximum, mean, median, and standard deviation) may not be available for every variable in the codebook. Conversely, a listing of frequencies in table format may not be present for every variable in the codebook either. However, all variables in the dataset are present and display sufficient information about each variable. These decisions are made intentionally and are at the discretion of the archive producing this codebook.

Center for Education and Drug Abuse Research (CEDAR): Etiological and Prospective Family Study in Southwestern Pennsylvania, Baseline Data, 1990-2011 [United States]

Psychiatric Treatment History Questionnaire - Father

ID SUBJECT ID

Location: 1-11 (width: 11; decimal: 0)
Variable Type: character
Based upon 715 valid cases out of 715 total cases.

ASSID Assessment ID

Location: 12-22 (width: 11; decimal: 0)
Variable Type: character
Based upon 715 valid cases out of 715 total cases.

VISIT Visit Number

Location: 23-24 (width: 2; decimal: 0)
Variable Type: numeric

<i>Value</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	715	100.0 %

Based upon 715 valid cases out of 715 total cases.

- Mean: 1.00
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 1.00
- Standard Deviation: 0.00

TRPC Had Treatment

Location: 25-25 (width: 1; decimal: 0)
Variable Type: numeric
Range of Missing Values (M): -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	415	58.0 %
1	Yes	300	42.0 %

Based upon 715 valid cases out of 715 total cases.

- Study 33444 -

- Mean: 0.42
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.49

TRPAE **Have you ever seen a professional because of emotional problems?**

Location: 26-26 (width: 1; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	526	73.6 %
1	Yes	189	26.4 %

Based upon 715 valid cases out of 715 total cases.

- Mean: 0.26
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.44

PCA21 **How old were you the first time you went for help because of emotional problems?**

Location: 27-28 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
5	-	2	0.3 %
6	-	1	0.1 %
9	-	2	0.3 %
10	-	5	0.7 %
11	-	1	0.1 %
12	-	2	0.3 %
13	-	4	0.6 %
14	-	5	0.7 %
15	-	1	0.1 %
16	-	1	0.1 %
17	-	5	0.7 %
18	-	3	0.4 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
19	-	3	0.4 %
20	-	2	0.3 %
21	-	3	0.4 %
22	-	4	0.6 %
23	-	1	0.1 %
24	-	7	1.0 %
25	-	5	0.7 %
26	-	4	0.6 %
27	-	3	0.4 %
28	-	4	0.6 %
29	-	3	0.4 %
30	-	9	1.3 %
31	-	8	1.1 %
32	-	6	0.8 %
33	-	9	1.3 %
34	-	13	1.8 %
35	-	9	1.3 %
36	-	5	0.7 %
37	-	12	1.7 %
38	-	6	0.8 %
39	-	5	0.7 %
40	-	7	1.0 %
41	-	5	0.7 %
42	-	5	0.7 %
43	-	3	0.4 %
44	-	3	0.4 %
45	-	3	0.4 %
46	-	2	0.3 %
47	-	1	0.1 %
48	-	1	0.1 %
50	-	3	0.4 %
54	-	1	0.1 %
55	-	1	0.1 %
56	-	1	0.1 %
-2 (M)	N/A	526	73.6 %

Based upon 189 valid cases out of 715 total cases.

- Mean: 30.49
- Median: 33.00
- Mode: 34.00

- Study 33444 -

- Minimum: 5.00
- Maximum: 56.00
- Standard Deviation: 10.69

PCA31

How old were you the last time you went for help because of emotional problems?

Location: 29-30 (width: 2; decimal: 0)

Variable Type: numeric

Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
9	-	1	0.1 %
10	-	2	0.3 %
11	-	1	0.1 %
12	-	2	0.3 %
13	-	4	0.6 %
14	-	3	0.4 %
15	-	1	0.1 %
18	-	2	0.3 %
20	-	1	0.1 %
21	-	3	0.4 %
22	-	2	0.3 %
23	-	1	0.1 %
24	-	3	0.4 %
25	-	4	0.6 %
26	-	5	0.7 %
27	-	5	0.7 %
28	-	4	0.6 %
29	-	5	0.7 %
30	-	6	0.8 %
31	-	4	0.6 %
32	-	5	0.7 %
33	-	8	1.1 %
34	-	8	1.1 %
35	-	7	1.0 %
36	-	7	1.0 %
37	-	11	1.5 %
38	-	9	1.3 %
39	-	10	1.4 %
40	-	8	1.1 %
41	-	9	1.3 %
42	-	7	1.0 %
43	-	6	0.8 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
44	-	9	1.3 %
45	-	5	0.7 %
46	-	6	0.8 %
47	-	2	0.3 %
48	-	1	0.1 %
50	-	2	0.3 %
51	-	2	0.3 %
52	-	2	0.3 %
53	-	2	0.3 %
54	-	1	0.1 %
56	-	1	0.1 %
58	-	2	0.3 %
-2 (M)	N/A	526	73.6 %

Based upon 189 valid cases out of 715 total cases.

- Mean: 34.96
- Median: 37.00
- Mode: 37.00
- Minimum: 9.00
- Maximum: 58.00
- Standard Deviation: 10.12

PCA41 How many times were you in treatment as an outpatient because of emotional problems?

Location: 31-32 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	127	17.8 %
2	-	23	3.2 %
3	-	14	2.0 %
4	-	5	0.7 %
5	-	2	0.3 %
6	-	1	0.1 %
7	-	1	0.1 %
10	-	1	0.1 %
24	-	1	0.1 %
50	-	1	0.1 %
90	-	1	0.1 %
-2 (M)	N/A	538	75.2 %

- Study 33444 -

Based upon 177 valid cases out of 715 total cases.

- Mean: 2.44
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 90.00
- Standard Deviation: 7.83

PCA51 **How many times were you in treatment as an inpatient because of emotional problems?**

Location: 33-34 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	30	4.2 %
2	-	12	1.7 %
3	-	8	1.1 %
5	-	1	0.1 %
6	-	2	0.3 %
7	-	1	0.1 %
14	-	1	0.1 %
37	-	1	0.1 %
-2 (M)	N/A	659	92.2 %

Based upon 56 valid cases out of 715 total cases.

- Mean: 2.73
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 37.00
- Standard Deviation: 5.12

TRPAG **Have you ever seen a professional because of drinking problems?**

Location: 35-35 (width: 1; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	608	85.0 %
1	Yes	107	15.0 %

Based upon 715 valid cases out of 715 total cases.

- Study 33444 -

- Mean: 0.15
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.36

PCA22

How old were you the first time you went for help because of drinking problems?

Location: 36-37 (width: 2; decimal: 0)

Variable Type: numeric

Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
16	-	3	0.4 %
17	-	3	0.4 %
19	-	2	0.3 %
20	-	1	0.1 %
21	-	1	0.1 %
22	-	3	0.4 %
23	-	6	0.8 %
24	-	3	0.4 %
25	-	5	0.7 %
26	-	8	1.1 %
27	-	2	0.3 %
28	-	5	0.7 %
29	-	1	0.1 %
30	-	5	0.7 %
31	-	4	0.6 %
32	-	6	0.8 %
33	-	7	1.0 %
34	-	3	0.4 %
35	-	6	0.8 %
36	-	5	0.7 %
37	-	7	1.0 %
38	-	5	0.7 %
39	-	4	0.6 %
40	-	3	0.4 %
41	-	1	0.1 %
42	-	3	0.4 %
43	-	1	0.1 %
44	-	2	0.3 %
50	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
51	-	1	0.1 %
-2 (M)	N/A	608	85.0 %

Based upon 107 valid cases out of 715 total cases.

- Mean: 31.06
- Median: 32.00
- Mode: 26.00
- Minimum: 16.00
- Maximum: 51.00
- Standard Deviation: 7.52

PCA32

How old were you the last time you went for help because of drinking problems?

Location: 38-39 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
17	-	2	0.3 %
19	-	1	0.1 %
21	-	1	0.1 %
23	-	2	0.3 %
24	-	1	0.1 %
25	-	2	0.3 %
26	-	1	0.1 %
27	-	2	0.3 %
28	-	3	0.4 %
29	-	5	0.7 %
30	-	3	0.4 %
31	-	3	0.4 %
32	-	6	0.8 %
33	-	7	1.0 %
34	-	6	0.8 %
35	-	8	1.1 %
36	-	7	1.0 %
37	-	4	0.6 %
38	-	6	0.8 %
39	-	8	1.1 %
40	-	5	0.7 %
41	-	3	0.4 %
42	-	6	0.8 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
43	-	2	0.3 %
44	-	5	0.7 %
45	-	2	0.3 %
47	-	1	0.1 %
51	-	3	0.4 %
53	-	1	0.1 %
54	-	1	0.1 %
-2 (M)	N/A	608	85.0 %

Based upon 107 valid cases out of 715 total cases.

- Mean: 35.62
- Median: 36.00
- Minimum: 17.00
- Maximum: 54.00
- Standard Deviation: 7.19

PCA42

How many times were you in treatment as an outpatient because of drinking problems?

Location: 40-41 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	57	8.0 %
2	-	15	2.1 %
3	-	6	0.8 %
5	-	1	0.1 %
7	-	1	0.1 %
10	-	1	0.1 %
50	-	2	0.3 %
-2 (M)	N/A	632	88.4 %

Based upon 83 valid cases out of 715 total cases.

- Mean: 2.73
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 50.00
- Standard Deviation: 7.59

PCA52

How many times were you in treatment as an inpatient because of drinking problems?

- Study 33444 -

Location: 42-43 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	47	6.6 %
2	-	14	2.0 %
3	-	6	0.8 %
4	-	7	1.0 %
5	-	3	0.4 %
6	-	3	0.4 %
7	-	1	0.1 %
10	-	1	0.1 %
12	-	1	0.1 %
15	-	1	0.1 %
-2 (M)	N/A	631	88.3 %

Based upon 84 valid cases out of 715 total cases.

- Mean: 2.36
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 15.00
- Standard Deviation: 2.46

TRPAR Have you ever seen a professional because of drug problems?

Location: 44-44 (width: 1; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	560	78.3 %
1	Yes	155	21.7 %

Based upon 715 valid cases out of 715 total cases.

- Mean: 0.22
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.41

PCA23 How old were you the first time you went for help because of drug problems?

- Study 33444 -

Location: 45-46 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
16	-	1	0.1 %
17	-	1	0.1 %
18	-	4	0.6 %
19	-	3	0.4 %
20	-	2	0.3 %
21	-	6	0.8 %
22	-	5	0.7 %
23	-	5	0.7 %
24	-	2	0.3 %
25	-	7	1.0 %
26	-	8	1.1 %
27	-	6	0.8 %
28	-	9	1.3 %
29	-	3	0.4 %
30	-	10	1.4 %
31	-	8	1.1 %
32	-	8	1.1 %
33	-	8	1.1 %
34	-	5	0.7 %
35	-	7	1.0 %
36	-	5	0.7 %
37	-	7	1.0 %
38	-	11	1.5 %
39	-	5	0.7 %
40	-	4	0.6 %
41	-	3	0.4 %
42	-	2	0.3 %
43	-	2	0.3 %
44	-	3	0.4 %
45	-	1	0.1 %
48	-	1	0.1 %
50	-	1	0.1 %
51	-	1	0.1 %
-2 (M)	N/A	560	78.3 %
-1 (M)	Missing, response refusal	1	0.1 %

Based upon 154 valid cases out of 715 total cases.

- Study 33444 -

- Mean: 31.16
- Median: 31.00
- Mode: 38.00
- Minimum: 16.00
- Maximum: 51.00
- Standard Deviation: 7.31

PCA33

How old were you the last time you went for help because of drug problems?

Location: 47-48 (width: 2; decimal: 0)

Variable Type: numeric

Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
17	-	1	0.1 %
18	-	1	0.1 %
21	-	2	0.3 %
22	-	1	0.1 %
23	-	1	0.1 %
24	-	1	0.1 %
25	-	3	0.4 %
26	-	2	0.3 %
27	-	2	0.3 %
28	-	3	0.4 %
29	-	4	0.6 %
30	-	5	0.7 %
31	-	5	0.7 %
32	-	7	1.0 %
33	-	11	1.5 %
34	-	7	1.0 %
35	-	11	1.5 %
36	-	8	1.1 %
37	-	8	1.1 %
38	-	11	1.5 %
39	-	7	1.0 %
40	-	10	1.4 %
41	-	8	1.1 %
42	-	9	1.3 %
43	-	5	0.7 %
44	-	7	1.0 %
45	-	6	0.8 %
46	-	1	0.1 %
47	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
48	-	1	0.1 %
51	-	4	0.6 %
54	-	1	0.1 %
-2 (M)	N/A	560	78.3 %
-1 (M)	Missing, response refusal	1	0.1 %

Based upon 154 valid cases out of 715 total cases.

- Mean: 36.46
- Median: 37.00
- Minimum: 17.00
- Maximum: 54.00
- Standard Deviation: 6.69

PCA43

How many times were you in treatment as an outpatient because of drug problems?

Location: 49-50 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	74	10.3 %
2	-	25	3.5 %
3	-	6	0.8 %
4	-	4	0.6 %
5	-	2	0.3 %
6	-	1	0.1 %
8	-	1	0.1 %
10	-	2	0.3 %
25	-	1	0.1 %
50	-	1	0.1 %
-2 (M)	N/A	598	83.6 %

Based upon 117 valid cases out of 715 total cases.

- Mean: 2.37
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 50.00
- Standard Deviation: 5.18

PCA53

How many times were you in treatment as an inpatient because of drug problems?

- Study 33444 -

Location: 51-52 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	70	9.8 %
2	-	22	3.1 %
3	-	14	2.0 %
4	-	7	1.0 %
5	-	7	1.0 %
6	-	4	0.6 %
7	-	1	0.1 %
8	-	1	0.1 %
10	-	2	0.3 %
12	-	1	0.1 %
15	-	1	0.1 %
16	-	1	0.1 %
30	-	1	0.1 %
-2 (M)	N/A	583	81.5 %

Based upon 132 valid cases out of 715 total cases.

- Mean: 2.66
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 30.00
- Standard Deviation: 3.52

TRPAO Thinking of the most difficult time when you saw a professional, was it as an Outpatient?

Location: 53-53 (width: 1; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	485	67.8 %
1	Yes	230	32.2 %

Based upon 715 valid cases out of 715 total cases.

- Mean: 0.32
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00

- Study 33444 -

- Standard Deviation: 0.47

PCA81 Whom did you see most frequently as an outpatient for Emotional Problems?

Location: 54-55 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Counselor	66	9.2 %
2	Social Worker	5	0.7 %
3	Psychologist	50	7.0 %
4	Psychiatrist	31	4.3 %
5	MD	11	1.5 %
6	Other	1	0.1 %
-2 (M)	N/A	551	77.1 %

Based upon 164 valid cases out of 715 total cases.

- Mean: 2.51
- Median: 3.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 6.00
- Standard Deviation: 1.39

PCA82 Whom did you see most frequently as an outpatient for Drinking Problems?

Location: 56-57 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Counselor	43	6.0 %
2	Social Worker	3	0.4 %
3	Psychologist	8	1.1 %
4	Psychiatrist	6	0.8 %
5	MD	1	0.1 %
6	Other	1	0.1 %
-2 (M)	N/A	653	91.3 %

Based upon 62 valid cases out of 715 total cases.

- Mean: 1.74
- Median: 1.00
- Mode: 1.00

- Study 33444 -

- Minimum: 1.00
- Maximum: 6.00
- Standard Deviation: 1.25

PCA83 Whom did you see most frequently as an outpatient for Drug Problems?

Location: 58-59 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Counselor	61	8.5 %
2	Social Worker	3	0.4 %
3	Psychologist	10	1.4 %
4	Psychiatrist	10	1.4 %
5	MD	5	0.7 %
6	Other	1	0.1 %
-2 (M)	N/A	625	87.4 %

Based upon 90 valid cases out of 715 total cases.

- Mean: 1.87
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 6.00
- Standard Deviation: 1.38

TRPAI Thinking of the most difficult time when you saw a professional, was it as an Inpatient?

Location: 60-60 (width: 1; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	551	77.1 %
1	Yes	164	22.9 %

Based upon 715 valid cases out of 715 total cases.

- Mean: 0.23
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.42

- Study 33444 -

PCA84 Whom did you see most frequently as an Inpatient for Emotional Problems?

Location: 61-62 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Counselor	8	1.1 %
2	Social Worker	1	0.1 %
3	Psychologist	2	0.3 %
4	Psychiatrist	33	4.6 %
5	MD	4	0.6 %
-2 (M)	N/A	667	93.3 %

Based upon 48 valid cases out of 715 total cases.

- Mean: 3.50
- Median: 4.00
- Mode: 4.00
- Minimum: 1.00
- Maximum: 5.00
- Standard Deviation: 1.22

PCA85 Whom did you see most frequently as an Inpatient for Drinking Problems?

Location: 63-64 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Counselor	51	7.1 %
2	Social Worker	1	0.1 %
3	Psychologist	8	1.1 %
4	Psychiatrist	12	1.7 %
5	MD	6	0.8 %
6	Other	3	0.4 %
-2 (M)	N/A	634	88.7 %

Based upon 81 valid cases out of 715 total cases.

- Mean: 2.14
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 6.00
- Standard Deviation: 1.61

- Study 33444 -

PCA86 Whom did you see most frequently as an Inpatient for Drug Problems?

Location: 65-66 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Counselor	79	11.0 %
2	Social Worker	2	0.3 %
3	Psychologist	12	1.7 %
4	Psychiatrist	16	2.2 %
5	MD	12	1.7 %
6	Other	3	0.4 %
-2 (M)	N/A	591	82.7 %

Based upon 124 valid cases out of 715 total cases.

- Mean: 2.10
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 6.00
- Standard Deviation: 1.59

PCOE9 How long did you go to this person as an Outpatient for Emotional Problems?

Location: 67-68 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Less than 1 month	33	4.6 %
2	1 to 3 months	42	5.9 %
3	4 to 6 months	16	2.2 %
4	7 to 12 months	32	4.5 %
5	13 months to 48 months	23	3.2 %
6	49 months or more	18	2.5 %
-2 (M)	N/A	551	77.1 %

Based upon 164 valid cases out of 715 total cases.

- Mean: 3.15
- Median: 3.00
- Mode: 2.00
- Minimum: 1.00
- Maximum: 6.00

- Study 33444 -

- Standard Deviation: 1.67

PCOG9 How long did you go to this person as an Outpatient for Drinking Problems?

Location: 69-70 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Less than 1 month	16	2.2 %
2	1 to 3 months	16	2.2 %
3	4 to 6 months	12	1.7 %
4	7 to 12 months	9	1.3 %
5	13 months to 48 months	5	0.7 %
6	49 months or more	4	0.6 %
-2 (M)	N/A	653	91.3 %

Based upon 62 valid cases out of 715 total cases.

- Mean: 2.73
- Median: 2.00
- Minimum: 1.00
- Maximum: 6.00
- Standard Deviation: 1.52

PCOR9 How long did you go to this person as an Outpatient for Drug Problems?

Location: 71-72 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Less than 1 month	23	3.2 %
2	1 to 3 months	22	3.1 %
3	4 to 6 months	16	2.2 %
4	7 to 12 months	10	1.4 %
5	13 months to 48 months	13	1.8 %
6	49 months or more	5	0.7 %
-2 (M)	N/A	626	87.6 %

Based upon 89 valid cases out of 715 total cases.

- Mean: 2.81
- Median: 2.00
- Mode: 1.00
- Minimum: 1.00

- Study 33444 -

- Maximum: 6.00
- Standard Deviation: 1.57

PCIE9 How long did you go to this person as an Inpatient for Emotional Problems?

Location: 73-74 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Less than 1 month	35	4.9 %
2	1 to 3 months	7	1.0 %
3	4 to 6 months	2	0.3 %
4	7 to 12 months	2	0.3 %
5	13 months to 48 months	2	0.3 %
-2 (M)	N/A	667	93.3 %

Based upon 48 valid cases out of 715 total cases.

- Mean: 1.52
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 5.00
- Standard Deviation: 1.05

PCIG9 How long did you go to this person as an Inpatient for Drinking Problems?

Location: 75-76 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Less than 1 month	53	7.4 %
2	1 to 3 months	19	2.7 %
3	4 to 6 months	4	0.6 %
4	7 to 12 months	4	0.6 %
5	13 months to 48 months	1	0.1 %
-2 (M)	N/A	634	88.7 %

Based upon 81 valid cases out of 715 total cases.

- Mean: 1.53
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00

- Study 33444 -

- Maximum: 5.00
- Standard Deviation: 0.90

PCIR9 How long did you go to this person as an Inpatient for Drug Problems?

Location: 77-78 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Less than 1 month	81	11.3 %
2	1 to 3 months	24	3.4 %
3	4 to 6 months	7	1.0 %
4	7 to 12 months	7	1.0 %
5	13 months to 48 months	2	0.3 %
6	49 months or more	3	0.4 %
-2 (M)	N/A	591	82.7 %

Based upon 124 valid cases out of 715 total cases.

- Mean: 1.66
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 6.00
- Standard Deviation: 1.17

PC101 What type of treatment did you have because of Emotional Problems - Outpatient Psychotherapy?

Location: 79-80 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	26	3.6 %
1	Yes	139	19.4 %
-2 (M)	N/A	550	76.9 %

Based upon 165 valid cases out of 715 total cases.

- Mean: 0.84
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.37

- Study 33444 -

PC105 **What type of treatment did you have because of Drinking Problems - Outpatient Psychotherapy?**

Location: 81-82 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	22	3.1 %
1	Yes	40	5.6 %
-2 (M)	N/A	653	91.3 %

Based upon 62 valid cases out of 715 total cases.

- Mean: 0.65
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.48

PC111 **What type of treatment did you have because of Drug Problems - Outpatient Psychotherapy?**

Location: 83-84 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	24	3.4 %
1	Yes	66	9.2 %
-2 (M)	N/A	625	87.4 %

Based upon 90 valid cases out of 715 total cases.

- Mean: 0.73
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.44

PC102 **What type of treatment did you have because of Emotional Problems - Outpatient ECT?**

Location: 85-86 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	162	22.7 %
1	Yes	2	0.3 %
-2 (M)	N/A	551	77.1 %

Based upon 164 valid cases out of 715 total cases.

- Mean: 0.01
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.11

PC106 **What type of treatment did you have because of Drinking Problems - Outpatient ECT?**

Location: 87-88 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	62	8.7 %
-2 (M)	N/A	653	91.3 %

Based upon 62 valid cases out of 715 total cases.

- Mean: 0.00
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 0.00
- Standard Deviation: 0.00

PC112 **What type of treatment did you have because of Drug Problems - Outpatient ECT?**

Location: 89-90 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	90	12.6 %
-2 (M)	N/A	625	87.4 %

Based upon 90 valid cases out of 715 total cases.

- Mean: 0.00

- Study 33444 -

- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 0.00
- Standard Deviation: 0.00

PC103 **What type of treatment did you have because of Emotional Problems - Outpatient Medication?**

Location: 91-92 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	124	17.3 %
1	Yes	40	5.6 %
-2 (M)	N/A	551	77.1 %

Based upon 164 valid cases out of 715 total cases.

- Mean: 0.24
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.43

PC107 **What type of treatment did you have because of Drinking Problems - Oupatient Medication?**

Location: 93-94 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	55	7.7 %
1	Yes	7	1.0 %
-2 (M)	N/A	653	91.3 %

Based upon 62 valid cases out of 715 total cases.

- Mean: 0.11
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.32

PC113 **What type of treatment did you have because of Drug Problems - Outpatient Medication?**

- Study 33444 -

Location: 95-96 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	72	10.1 %
1	Yes	18	2.5 %
-2 (M)	N/A	625	87.4 %

Based upon 90 valid cases out of 715 total cases.

- Mean: 0.20
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.40

PC108 **What type of treatment did you have because of Drinking Problems - Outpatient Detoxification?**

Location: 97-98 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	58	8.1 %
1	Yes	4	0.6 %
-2 (M)	N/A	653	91.3 %

Based upon 62 valid cases out of 715 total cases.

- Mean: 0.06
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.25

PC114 **What type of treatment did you have because of Drug Problems - Outpatient Detoxification?**

Location: 99-100 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	82	11.5 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Yes	8	1.1 %
-2 (M)	N/A	625	87.4 %

Based upon 90 valid cases out of 715 total cases.

- Mean: 0.09
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.29

PC109 **What type of treatment did you have because of Drinking Problems - Outpatient Rehabilitation?**

Location: 101-102 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	38	5.3 %
1	Yes	24	3.4 %
-2 (M)	N/A	653	91.3 %

Based upon 62 valid cases out of 715 total cases.

- Mean: 0.39
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.49

PC115 **What type of treatment did you have because of Drug Problems - Outpatient Rehabilitation?**

Location: 103-104 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	63	8.8 %
1	Yes	27	3.8 %
-2 (M)	N/A	625	87.4 %

Based upon 90 valid cases out of 715 total cases.

- Study 33444 -

- Mean: 0.30
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.46

PC104 **What type of treatment did you have because of Emotional Problems - Other Outpatient?**

Location: 105-106 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	157	22.0 %
1	Yes	6	0.8 %
-2 (M)	N/A	552	77.2 %

Based upon 163 valid cases out of 715 total cases.

- Mean: 0.04
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.19

PC110 **What type of treatment did you have because of Drinking Problems - Other Outpatient**

Location: 107-108 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	59	8.3 %
1	Yes	3	0.4 %
-2 (M)	N/A	653	91.3 %

Based upon 62 valid cases out of 715 total cases.

- Mean: 0.05
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.22

- Study 33444 -

PC116 **What type of treatment did you have because of Drug Problems - Other Outpatient**

Location: 109-110 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	85	11.9 %
1	Yes	5	0.7 %
-2 (M)	N/A	625	87.4 %

Based upon 90 valid cases out of 715 total cases.

- Mean: 0.06
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.23

PC117 **What type of treatment did you have because of Emotional Problems - Inpatient Psychotherapy?**

Location: 111-112 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	10	1.4 %
1	Yes	41	5.7 %
-2 (M)	N/A	664	92.9 %

Based upon 51 valid cases out of 715 total cases.

- Mean: 0.80
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.40

PC123 **What type of treatment did you have because of Drinking Problems - Inpatient Psychotherapy?**

Location: 113-114 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	29	4.1 %
1	Yes	52	7.3 %
-2 (M)	N/A	634	88.7 %

Based upon 81 valid cases out of 715 total cases.

- Mean: 0.64
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.48

PC129 **What type of treatment did you have because of Drug Problems - Inpatient Psychotherapy?**

Location: 115-116 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	39	5.5 %
1	Yes	84	11.7 %
-2 (M)	N/A	592	82.8 %

Based upon 123 valid cases out of 715 total cases.

- Mean: 0.68
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.47

PC118 **What type of treatment did you have because of Emotional Problems - Inpatient ECT?**

Location: 117-118 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	51	7.1 %
-2 (M)	N/A	664	92.9 %

Based upon 51 valid cases out of 715 total cases.

- Study 33444 -

- Mean: 0.00
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 0.00
- Standard Deviation: 0.00

PC124 **What type of treatment did you have because of Drinking Problems - Inpatient ECT?**

Location: 119-120 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	81	11.3 %
-2 (M)	N/A	634	88.7 %

Based upon 81 valid cases out of 715 total cases.

- Mean: 0.00
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 0.00
- Standard Deviation: 0.00

PC130 **What type of treatment did you have because of Drug Problems - Inpatient ECT?**

Location: 121-122 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	123	17.2 %
-2 (M)	N/A	592	82.8 %

Based upon 123 valid cases out of 715 total cases.

- Mean: 0.00
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 0.00
- Standard Deviation: 0.00

PC119 **What type of treatment did you have because of Emotional Problems - Inpatient Medication?**

- Study 33444 -

Location: 123-124 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	22	3.1 %
1	Yes	29	4.1 %
-2 (M)	N/A	664	92.9 %

Based upon 51 valid cases out of 715 total cases.

- Mean: 0.57
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.50

PC125 What type of treatment did you have because of Drinking Problems - Inpatient Medication?

Location: 125-126 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	60	8.4 %
1	Yes	21	2.9 %
-2 (M)	N/A	634	88.7 %

Based upon 81 valid cases out of 715 total cases.

- Mean: 0.26
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.44

PC131 What type of treatment did you have because of Drug Problems - Inpatient Medication?

Location: 127-128 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	88	12.3 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	Yes	35	4.9 %
-2 (M)	N/A	592	82.8 %

Based upon 123 valid cases out of 715 total cases.

- Mean: 0.28
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.45

PC120 **What type of treatment did you have because of Emotional Problems - Inpatient Detoxification?**

Location: 129-130 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	50	7.0 %
1	Yes	1	0.1 %
-2 (M)	N/A	664	92.9 %

Based upon 51 valid cases out of 715 total cases.

- Mean: 0.02
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.14

PC126 **What type of treatment did you have because of Drinking Problems - Inpatient Detoxification?**

Location: 131-132 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	25	3.5 %
1	Yes	56	7.8 %
-2 (M)	N/A	634	88.7 %

Based upon 81 valid cases out of 715 total cases.

- Study 33444 -

- Mean: 0.69
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.46

PC132 **What type of treatment did you have because of Drug Problems - Inpatient Detoxification?**

Location: 133-134 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	39	5.5 %
1	Yes	84	11.7 %
-2 (M)	N/A	592	82.8 %

Based upon 123 valid cases out of 715 total cases.

- Mean: 0.68
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.47

PC121 **What type of treatment did you have because of Emotional Problems - Inpatient Rehabilitation?**

Location: 135-136 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	51	7.1 %
-2 (M)	N/A	664	92.9 %

Based upon 51 valid cases out of 715 total cases.

- Mean: 0.00
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 0.00
- Standard Deviation: 0.00

- Study 33444 -

PC127 What type of treatment did you have because of Drinking Problems - Inpatient Rehabilitation?

Location: 137-138 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	36	5.0 %
1	Yes	45	6.3 %
-2 (M)	N/A	634	88.7 %

Based upon 81 valid cases out of 715 total cases.

- Mean: 0.56
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.50

PC133 What type of treatment did you have because of Drug Problems - Inpatient Rehabilitation?

Location: 139-140 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	54	7.6 %
1	Yes	70	9.8 %
-2 (M)	N/A	591	82.7 %

Based upon 124 valid cases out of 715 total cases.

- Mean: 0.56
- Median: 1.00
- Mode: 1.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.50

PC122 What type of treatment did you have because of Emotional Problems - Other Inpatient?

Location: 141-142 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	48	6.7 %
1	Yes	3	0.4 %
-2 (M)	N/A	664	92.9 %

Based upon 51 valid cases out of 715 total cases.

- Mean: 0.06
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.24

PC128

What type of treatment did you have because of Drinking Problems - Other Inpatient?

Location: 143-144 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	78	10.9 %
1	Yes	3	0.4 %
-2 (M)	N/A	634	88.7 %

Based upon 81 valid cases out of 715 total cases.

- Mean: 0.04
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.19

PC134

What type of treatment did you have because of Drug Problems - Other Inpatient?

Location: 145-146 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	117	16.4 %
1	Yes	5	0.7 %
-2 (M)	N/A	593	82.9 %

Based upon 122 valid cases out of 715 total cases.

- Study 33444 -

- Mean: 0.04
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.20

PCB1 Self-help Groups: 1. Alcoholics Anonymous (AA)

Location: 147-148 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	615	86.0 %
1	Yes	99	13.8 %
-2 (M)	N/A	1	0.1 %

Based upon 714 valid cases out of 715 total cases.

- Mean: 0.14
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.35

PCB2 Self-help Groups: 2. Narcotics Anonymous (NA)

Location: 149-150 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	607	84.9 %
1	Yes	107	15.0 %
-2 (M)	N/A	1	0.1 %

Based upon 714 valid cases out of 715 total cases.

- Mean: 0.15
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.36

PCB3 Self-help Groups: 3. Cocaine Anonymous (CA)

Location: 151-152 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	706	98.7 %
1	Yes	8	1.1 %
-2 (M)	N/A	1	0.1 %

Based upon 714 valid cases out of 715 total cases.

- Mean: 0.01
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.11

TRIIIE Untreated Psychopathology: 1) Were there any (other) times when you or someone else felt that you needed help because of emotional problems and you did not get it?

Location: 153-153 (width: 1; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	641	89.7 %
1	Yes	74	10.3 %

Based upon 715 valid cases out of 715 total cases.

- Mean: 0.10
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.30

TRIIG Untreated Psychopathology: 1) Were there any (other) times when you or someone else felt that you needed help because of drinking problems and you did not get it?

Location: 154-154 (width: 1; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	656	91.7 %
1	Yes	59	8.3 %

Based upon 715 valid cases out of 715 total cases.

- Mean: 0.08
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.28

TRIIR Untreated Psychopathology: 1) Were there any (other) times when you or someone else felt that you needed help because of drug problems and you did not get it?

Location: 155-155 (width: 1; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
0	No	640	89.5 %
1	Yes	75	10.5 %

Based upon 715 valid cases out of 715 total cases.

- Mean: 0.10
- Median: 0.00
- Mode: 0.00
- Minimum: 0.00
- Maximum: 1.00
- Standard Deviation: 0.31

PCU02 Worst time you needed professional help because of emotional problems and you did not get it: 2) What was the reason?

Location: 156-165 (width: 10; decimal: 0)
 Variable Type: character
 Range of Missing Values (M): -1 , -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
-1 (M)	Missing, response refusal	6	0.8 %
-2 (M)	N/A	641	89.7 %

Based upon 68 valid cases out of 715 total cases.

- Study 33444 -

PCU03 Worst time you needed professional help because of emotional problems and you did not get it: 3) How old were you when you first needed help?

Location: 166-167 (width: 2; decimal: 0)

Variable Type: numeric

Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
6	-	2	0.3 %
7	-	1	0.1 %
8	-	1	0.1 %
10	-	1	0.1 %
11	-	1	0.1 %
12	-	1	0.1 %
14	-	2	0.3 %
16	-	2	0.3 %
17	-	1	0.1 %
18	-	3	0.4 %
19	-	1	0.1 %
20	-	3	0.4 %
21	-	2	0.3 %
22	-	4	0.6 %
23	-	4	0.6 %
24	-	2	0.3 %
25	-	2	0.3 %
26	-	1	0.1 %
27	-	3	0.4 %
28	-	2	0.3 %
29	-	2	0.3 %
30	-	4	0.6 %
31	-	3	0.4 %
32	-	2	0.3 %
34	-	4	0.6 %
35	-	2	0.3 %
37	-	4	0.6 %
38	-	1	0.1 %
39	-	1	0.1 %
40	-	2	0.3 %
41	-	1	0.1 %
43	-	1	0.1 %
44	-	1	0.1 %
45	-	1	0.1 %
46	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
50	-	1	0.1 %
54	-	2	0.3 %
-2 (M)	N/A	641	89.7 %
-1 (M)	Missing, response refusal	2	0.3 %

Based upon 72 valid cases out of 715 total cases.

- Mean: 27.44
- Median: 27.00
- Minimum: 6.00
- Maximum: 54.00
- Standard Deviation: 11.06

PCU04 Worst time you needed professional help because of emotional problems and you did not get it: 4) How old were you the last time you needed help?

Location: 168-169 (width: 2; decimal: 0)

Variable Type: numeric

Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
10	-	1	0.1 %
11	-	1	0.1 %
12	-	2	0.3 %
16	-	1	0.1 %
17	-	1	0.1 %
19	-	1	0.1 %
20	-	1	0.1 %
22	-	5	0.7 %
23	-	4	0.6 %
25	-	3	0.4 %
26	-	1	0.1 %
27	-	3	0.4 %
28	-	2	0.3 %
29	-	2	0.3 %
30	-	3	0.4 %
31	-	2	0.3 %
32	-	3	0.4 %
33	-	1	0.1 %
34	-	3	0.4 %
35	-	4	0.6 %
36	-	2	0.3 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
37	-	1	0.1 %
38	-	4	0.6 %
39	-	4	0.6 %
41	-	1	0.1 %
42	-	2	0.3 %
43	-	3	0.4 %
45	-	5	0.7 %
46	-	1	0.1 %
49	-	1	0.1 %
50	-	1	0.1 %
51	-	1	0.1 %
54	-	2	0.3 %
60	-	1	0.1 %
-2 (M)	N/A	641	89.7 %
-1 (M)	Missing, response refusal	1	0.1 %

Based upon 73 valid cases out of 715 total cases.

- Mean: 32.85
- Median: 33.00
- Minimum: 10.00
- Maximum: 60.00
- Standard Deviation: 10.85

PCU05

Worst time you needed professional help because of emotional problems and you did not get it: 5) How many times did you feel you needed help?

Location: 170-171 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	50	7.0 %
2	-	4	0.6 %
3	-	2	0.3 %
5	-	3	0.4 %
7	-	1	0.1 %
8	-	1	0.1 %
10	-	2	0.3 %
12	-	1	0.1 %
15	-	1	0.1 %
20	-	2	0.3 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
50	-	1	0.1 %
99	-	1	0.1 %
-2 (M)	N/A	642	89.8 %
-1 (M)	Missing, response refusal	4	0.6 %

Based upon 69 valid cases out of 715 total cases.

- Mean: 4.78
- Median: 1.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 99.00
- Standard Deviation: 13.48

PCU06 Worst time you needed professional help because of emotional problems and you did not get it: 6) During this period for how long (# days) did you need help but not get it

Location: 172-174 (width: 3; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
14	-	1	0.1 %
20	-	1	0.1 %
21	-	1	0.1 %
30	-	7	1.0 %
36	-	1	0.1 %
42	-	1	0.1 %
60	-	4	0.6 %
65	-	1	0.1 %
72	-	2	0.3 %
90	-	3	0.4 %
120	-	1	0.1 %
180	-	1	0.1 %
182	-	1	0.1 %
240	-	1	0.1 %
300	-	2	0.3 %
365	-	7	1.0 %
375	-	1	0.1 %
540	-	1	0.1 %
548	-	1	0.1 %
730	-	6	0.8 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
780	-	1	0.1 %
999	-	27	3.8 %
-2 (M)	N/A	642	89.8 %
-1 (M)	Missing, response refusal	1	0.1 %

Based upon 72 valid cases out of 715 total cases.

- Mean: 535.21
- Median: 457.50
- Mode: 999.00
- Minimum: 14.00
- Maximum: 999.00
- Standard Deviation: 414.08

PCU07 Worst time you needed professional help because of drinking problems and you did not get it: 2) What was the reason?

Location: 175-184 (width: 10; decimal: 0)
 Variable Type: character
 Range of Missing Values (M): -1 , -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
-1 (M)	Missing, response refusal	5	0.7 %
-2 (M)	N/A	656	91.7 %

Based upon 54 valid cases out of 715 total cases.

PCU08 Worst time you needed professional help because of drinking problems and you did not get it: 3) How old were you when you first needed help?

Location: 185-186 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
11	-	1	0.1 %
13	-	1	0.1 %
14	-	1	0.1 %
16	-	1	0.1 %
17	-	4	0.6 %
18	-	4	0.6 %
19	-	2	0.3 %
20	-	4	0.6 %
21	-	4	0.6 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
23	-	1	0.1 %
24	-	1	0.1 %
25	-	3	0.4 %
26	-	1	0.1 %
27	-	3	0.4 %
28	-	4	0.6 %
29	-	2	0.3 %
30	-	9	1.3 %
31	-	1	0.1 %
32	-	1	0.1 %
34	-	1	0.1 %
35	-	4	0.6 %
36	-	1	0.1 %
37	-	2	0.3 %
38	-	1	0.1 %
39	-	1	0.1 %
54	-	1	0.1 %
-2 (M)	N/A	656	91.7 %

Based upon 59 valid cases out of 715 total cases.

- Mean: 26.15
- Median: 27.00
- Mode: 30.00
- Minimum: 11.00
- Maximum: 54.00
- Standard Deviation: 7.93

PCU09 Worst time you needed professional help because of drinking problems and you did not get it: 4) How old were you the last time you needed help?

Location: 187-188 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
18	-	1	0.1 %
20	-	1	0.1 %
22	-	2	0.3 %
24	-	1	0.1 %
25	-	3	0.4 %
26	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
27	-	1	0.1 %
28	-	3	0.4 %
29	-	5	0.7 %
30	-	1	0.1 %
31	-	3	0.4 %
32	-	4	0.6 %
33	-	2	0.3 %
34	-	2	0.3 %
35	-	4	0.6 %
36	-	5	0.7 %
37	-	2	0.3 %
38	-	6	0.8 %
39	-	2	0.3 %
40	-	2	0.3 %
41	-	1	0.1 %
43	-	1	0.1 %
45	-	1	0.1 %
50	-	1	0.1 %
51	-	1	0.1 %
52	-	1	0.1 %
54	-	2	0.3 %
-2 (M)	N/A	656	91.7 %

Based upon 59 valid cases out of 715 total cases.

- Mean: 34.20
- Median: 34.00
- Mode: 38.00
- Minimum: 18.00
- Maximum: 54.00
- Standard Deviation: 7.95

PCU10

Worst time you needed professional help because of drinking problems and you did not get it: 5) How many times did you feel you needed help?

Location: 189-190 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	27	3.8 %
2	-	3	0.4 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
3	-	4	0.6 %
4	-	1	0.1 %
5	-	1	0.1 %
7	-	1	0.1 %
10	-	3	0.4 %
12	-	3	0.4 %
14	-	2	0.3 %
15	-	1	0.1 %
20	-	3	0.4 %
25	-	1	0.1 %
30	-	1	0.1 %
50	-	2	0.3 %
99	-	3	0.4 %
-2 (M)	N/A	656	91.7 %
-1 (M)	Missing, response refusal	3	0.4 %

Based upon 56 valid cases out of 715 total cases.

- Mean: 12.18
- Median: 2.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 99.00
- Standard Deviation: 23.49

PCU11 Worst time you needed professional help because of drinking problems and you did not get it: 6) During this period for how long (# days) did you need help but not get it?

Location: 191-193 (width: 3; decimal: 0)

Variable Type: numeric

Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
14	-	1	0.1 %
30	-	2	0.3 %
60	-	1	0.1 %
84	-	1	0.1 %
112	-	1	0.1 %
120	-	1	0.1 %
150	-	1	0.1 %
365	-	9	1.3 %
375	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
450	-	1	0.1 %
548	-	1	0.1 %
700	-	1	0.1 %
730	-	3	0.4 %
760	-	1	0.1 %
780	-	1	0.1 %
999	-	32	4.5 %
-2 (M)	N/A	656	91.7 %
-1 (M)	Missing, response refusal	1	0.1 %

Based upon 58 valid cases out of 715 total cases.

- Mean: 718.21
- Median: 999.00
- Mode: 999.00
- Minimum: 14.00
- Maximum: 999.00
- Standard Deviation: 355.35

PCU12 Worst time you needed professional help because of drug problems and you did not get it: 2) What was the reason?

Location: 194-203 (width: 10; decimal: 0)
 Variable Type: character
 Range of Missing Values (M): -1 , -2

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
-1 (M)	Missing, response refusal	7	1.0 %
-2 (M)	N/A	640	89.5 %

Based upon 68 valid cases out of 715 total cases.

PCU14 Worst time you needed professional help because of drug problems and you did not get it: 4) How old were you when you last needed help?

Location: 204-205 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
17	-	2	0.3 %
18	-	2	0.3 %
22	-	1	0.1 %
23	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
24	-	2	0.3 %
25	-	3	0.4 %
26	-	3	0.4 %
27	-	3	0.4 %
28	-	2	0.3 %
29	-	5	0.7 %
30	-	1	0.1 %
31	-	4	0.6 %
32	-	4	0.6 %
33	-	4	0.6 %
34	-	2	0.3 %
35	-	5	0.7 %
36	-	4	0.6 %
37	-	2	0.3 %
38	-	8	1.1 %
39	-	2	0.3 %
40	-	3	0.4 %
41	-	2	0.3 %
42	-	1	0.1 %
43	-	1	0.1 %
44	-	2	0.3 %
45	-	1	0.1 %
48	-	1	0.1 %
51	-	2	0.3 %
54	-	1	0.1 %
-2 (M)	N/A	640	89.5 %
-1 (M)	Missing, response refusal	1	0.1 %

Based upon 74 valid cases out of 715 total cases.

- Mean: 33.50
- Median: 33.50
- Mode: 38.00
- Minimum: 17.00
- Maximum: 54.00
- Standard Deviation: 7.81

PCU13

**Worst time you needed professional help because of drug problems and you did not get it:
3) How old were you the first time you needed help?**

Location: 206-207 (width: 2; decimal: 0)

Variable Type: numeric

- Study 33444 -

Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
13	-	1	0.1 %
16	-	2	0.3 %
17	-	4	0.6 %
18	-	5	0.7 %
19	-	2	0.3 %
20	-	2	0.3 %
21	-	3	0.4 %
22	-	2	0.3 %
23	-	1	0.1 %
24	-	3	0.4 %
25	-	6	0.8 %
26	-	1	0.1 %
27	-	2	0.3 %
28	-	8	1.1 %
29	-	2	0.3 %
30	-	9	1.3 %
31	-	2	0.3 %
32	-	3	0.4 %
33	-	1	0.1 %
34	-	2	0.3 %
35	-	2	0.3 %
36	-	3	0.4 %
37	-	1	0.1 %
38	-	1	0.1 %
39	-	1	0.1 %
40	-	2	0.3 %
50	-	1	0.1 %
52	-	1	0.1 %
54	-	1	0.1 %
-2 (M)	N/A	640	89.5 %
-1 (M)	Missing, response refusal	1	0.1 %

Based upon 74 valid cases out of 715 total cases.

- Mean: 27.73
- Median: 28.00
- Mode: 30.00
- Minimum: 13.00
- Maximum: 54.00
- Standard Deviation: 8.24

- Study 33444 -

PCU15 Worst time you needed professional help because of drug problems and you did not get it: 5) How many times did you feel you needed help?

Location: 208-209 (width: 2; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
1	-	33	4.6 %
2	-	7	1.0 %
3	-	3	0.4 %
4	-	1	0.1 %
5	-	1	0.1 %
6	-	1	0.1 %
7	-	1	0.1 %
10	-	5	0.7 %
12	-	3	0.4 %
14	-	2	0.3 %
15	-	3	0.4 %
20	-	3	0.4 %
25	-	2	0.3 %
42	-	1	0.1 %
50	-	2	0.3 %
99	-	4	0.6 %
-2 (M)	N/A	641	89.7 %
-1 (M)	Missing, response refusal	2	0.3 %

Based upon 72 valid cases out of 715 total cases.

- Mean: 12.29
- Median: 2.00
- Mode: 1.00
- Minimum: 1.00
- Maximum: 99.00
- Standard Deviation: 23.66

PCU16 Worst time you needed professional help because of drug problems and you did not get it: 6) During this period for how long (# days) did you need help but not get it?

Location: 210-212 (width: 3; decimal: 0)
 Variable Type: numeric
 Range of Missing Values (M): -2 , -1

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
14	-	1	0.1 %

- Study 33444 -

<i>Value</i>	<i>Label</i>	<i>Unweighted Frequency</i>	<i>%</i>
30	-	2	0.3 %
60	-	1	0.1 %
84	-	1	0.1 %
90	-	1	0.1 %
100	-	1	0.1 %
120	-	1	0.1 %
150	-	2	0.3 %
270	-	1	0.1 %
300	-	1	0.1 %
365	-	11	1.5 %
375	-	1	0.1 %
500	-	1	0.1 %
600	-	1	0.1 %
630	-	1	0.1 %
730	-	6	0.8 %
760	-	1	0.1 %
999	-	41	5.7 %
-2 (M)	N/A	640	89.5 %

Based upon 75 valid cases out of 715 total cases.

- Mean: 714.89
- Median: 999.00
- Mode: 999.00
- Minimum: 14.00
- Maximum: 999.00
- Standard Deviation: 353.15